

A Spectroscopic Study of Vega (α Lyrae) and Sirius (α Canis Majoris)

Kenneth R Whight 01/11/2022

Abstract

This paper describes an investigation into what can be learned about the physical properties of the White star Vega (α Lyrae) from both low (150 lines/mm) and high (2400 lines/mm) resolution spectra, based on the simple model that the star is a rotating uniformly emitting oblate spheroid with a photosphere that is a single layer in thermal equilibrium. High resolution spectra of Sirius (α Canis Majoris) are remarkably similar to those obtained for Vega so the results of studying Vega are assumed to apply equally well to Sirius. Both stars are hot A0 V stars, Vega is known to be rotating rapidly and we are viewing it from within 5° of its pole so the effects of rotation are assumed negligible. The aim of this work was to test the ability of a simple stellar model to predict the Hydrogen line profiles in Vega's and therefore also Sirius' spectrum.

1. Introduction

Vega (α Lyr) is classed as a A0 V star i.e. a hot (A0) main sequence (V) star that is rotating rapidly but is being viewed from within 5° of its pole so that the effects of rotation on its spectra should be negligible. Sirius (α CMa) is a similarly classed star whose spectra are remarkably similar to those of Vega, however it is not recorded as a fast rotator. I shall use the similarity of the spectra to assert that the results of modelling Vega apply equally well to Sirius.

The aim of this work was to test the ability of a simple stellar model to predict the Hydrogen line profiles in Vega's and therefore by implication, Sirius' spectrum.


The stellar model used was that of a, solid body, rotating uniformly emitting oblate spheroid with a photosphere that is a single layer in thermal equilibrium. It is also assumed that the observed absorption lines are formed solely within this photosphere.

Using this model an effective "black body" temperature can be deduced from low resolution (150 lines/mm) spectra provided proper calibration is performed to correct the continuum spectrum for instrument response and atmospheric absorption. High resolution (2400 lines/mm) investigations of individual line shapes can then be used to determine other model parameters for example, the "Lorentzian Half Width" of pressure induced line broadening and, if appropriate, the star's speed of rotation.


The theory, methods and computer programs used in this study have been previously described in earlier studies of the blue component of Albireo (β Cyg) and Deneb (α Cyg) that are published on my website www.thewhightstuff.co.uk.

2 Low Resolution Spectra

Figure 1 shows a low resolution (150 lines/mm) spectrum of Vega, this spectrum was fully calibrated for instrument response and atmospheric absorption using a library reference spectrum. In the figure the Hydrogen α , β and γ line positions have been indicated.






Low resolution data can be used to obtain an estimate for the effective temperature of a star. It is simply necessary to divide the spectrum by the particular "Planck wavelength curve" that results in the flattest resultant spectrum. Figure 2 shows the optimal flattened spectrum of Vega after division by a 14025K Planck curve.



However it is worth noting that significant errors can result for hot stars as the peak of emission is well into the UV region of the spectrum so the resulting temperature is very sensitive to small "errors" in the measured spectrum. Luckily line profiles are most sensitive to pressure effects and not temperature. I have used the temperature suggested by the low resolution measurements in my simulations.

3.0 High Resolution Spectra

High resolution (2400 line/mm) spectra, of both Vega and Sirius, taken at H_γ , H_β and H_α wavelengths are shown in figures 3, 4 and 5 respectively, apart from small differences in position, due to Doppler effects, or experimental error, they are remarkably similar.




All lines appear to show evidence of a "spike" that seems to emerge out of the broader body of the line as we progress from H_γ to H_α via H_β . The line shapes are not as "rounded" as seen in my earlier studies of AlbireoB and Deneb. There is the possibility that this could be due to a non-uniformity

in the emission from the star (www.star-facts.com) or the result of the photosphere having two distinct layers. These possibilities will be investigate in a subsequent paper.

In this paper I will model Vega on the same basis as my work on Albireo B and Deneb, i.e. having a photosphere that is a single layer in thermal equilibrium.


3.1 Single Layer Model

For the H_β absorption line the central wavelength was determined, based on equal areas each side of centre, to be 4861.47A. The absorption profile was then modelled assuming the absorption line amplitude at the line centre $A_\beta(\lambda_\beta) = 0.204$, the result of which is shown in figure 6.



In this figure it can be seen that a compromise has had to be made between fitting the shape of the "core" and the "wings" of the profile. Important parameter values appear as labels in the RSpec displayed spectra.

Subsequent prediction of the H_γ and H_α absorption lines, using the method described in my previous reports yield the results depicted in figures 7 and 8.



In these figures it can be seen that the model predictions are not too far off from the measurements with regard to the amplitude of each line however there is an obvious difference in shape between the modelled and measured profiles. It is to be hoped that, as for the Sun, although the line profile shapes were not very well reproduced the calculations

provide a reasonable estimate for the thickness and average pressure of Vega's photosphere.





Figure 9 shows the result of calculations of the thickness and pressure of Vega's photosphere as a function of an, as yet to be determined, impact parameter.



- The relative amplitude of the hydrogen Balmer line series are well reproduced by the simple model however the line profiles are not well modelled by a uniform single layer photosphere.
- The temperature of the stars photospheres are 14025K.
- A future study that assumes the photosphere comprises two regions in parallel or two layers in series (or combination thereof) may yield a better fit to the spectral line profiles.

4.0 Discussion

The single layer model has proven reasonably successful in modelling other stars e.g. the Sun, Albireo B and Deneb, with regard to both amplitude and line profile shape when known complicating factors (Albireo B's decretion disc and Deneb's outflows) are allowed for. However for Vega, whilst line amplitudes are well reproduced the shape of the line profiles are not well reproduced. It is not obvious what process could yield such line shapes but failure of the uniformity of emission or the assumption of a single layer are the obvious candidates.

I will investigate further in the near future to see if two layer/region models can better reproduce the spectral line profiles obtained from Vega.

5.0 Conclusions

A spectroscopic study of Vega (α Lyrae) and Sirius (α Canis Majoris) has been performed to determine physical properties of these stars. It has been found that:-